UpFront

423 Highlights from this issue R M Beattie

Commentary

425 Training in endoscopic haemostasis: targeting the bleeding point I D Penman, N I Church

Leading article

427 Infliximab in acute severe colitis: getting the right dose J Fiske, T Conley, S Sebastian, S Subramanian

Original research

- **430** Training in endotherapy for acute upper gastrointestinal bleeding: a UK-wide gastroenterology trainee survey J Segal, K Siau, C Kanagasundaram, A Askari, P Dunckley, A J Morris
- **436** Variation in exposure to endoscopic haemostasis for acute upper gastrointestinal bleeding during UK gastroenterology training K Siau, A J Morris, A Murugananthan, B McKaig, P Dunckley
- **441** Comparison of medium to longterm outcomes of acute severe ulcerative colitis patients receiving accelerated and standard infliximab induction D J Gibson, J Doherty, M McNally,

J Campion, D Keegan, A Keogh,

- U Kennedy, K Byrne, L J Egan,
- S McKiernan, FI MacCarthy, S Sengupta,
- J Sheridan, H E Mulcahy, G Cullen,
- E Slattery, D Kevans, G A Doherty

- 448 Measuring patient and carer experience related to paediatric gastrointestinal endoscopy: multicentre questionnaire study I W Y Wan, E Griffiths, R Rabone, Z Zuhair, Z Londt, D Rawat, J Maginnis, A Elzubair, A Pigott, M Thomson, P Narula
- 454 Impact of the COVID-19 outbreak on endoscopy training in a tertiary care centre in Germany C Roemmele, J Manzeneder, H Messmann, A Ebigbo

Reviews

- 458 Acute-on-chronic liver failure: definition, prognosis and management A Amin, R P Mookerjee
- 468 How to manage IBD in the 'elderly' J P Segal, H M T Htet, J Limdi, BH Hayee
- 478 Non-alcoholic fatty liver disease (NAFLD) in non-obese individuals M Phipps, J Wattacheril

Education in practice

484 Diagnostic ileocolonoscopy: getting the basics right M C Choy, M Matharoo, S Thomas-Gibson



Cover image Colonic polyp, double contrast coloured barium x-ray. SOVEREIGN, ISM/ SCIENCE PHOTO LIBRARY.






This article has been chosen by the Editor to be of special interest

or importance and is freely available online.

This article has been made freely available online under the BMJ Journals open access scheme. OPEN ACCESS

See http://fg.bmj.com/site/about/guidelines.xhtml#open



www.publicationethics.org.uk




C O P E

Member since 2009

 \checkmark

6



Opinion

491 Number of significant polyps detected per six-minute withdrawal time at colonoscopy (SP6): a new measure of colonoscopy efficiency and quality R Rameshshanker, B P Saunders

Letters

- **494** Practical plus personal: a refreshing approach to careers advice by the British Society of Gastroenterology Taster Course C Boyd
- **494** Faecal incontinence is not rare in irritable bowel syndrome C J Black, A C Ford

496 Endoscopy training in upper GI bleeding for UK trainees in Khartoum, Sudan: South Yorkshire experience A Al-Rifaie, T Min, D Allam, K Smith, D P Hurlstone, A Abdo, E Said, M Donnelly

Highlights from the literature

498 Highlights from the literature R M Beattie

Inside view

- **499** Unusual radiological sign in a gentleman with recurrent dysphagia M Y W Wong, M Richards, B Holt, G S Hebbard
- 501 Mass in a fatty liver D J van der Windt, G C Wilson, I A Ziogas, S Tohme, A Furlan, M A Nalesnik, D A Geller