Supplementary material Frontline Gastroenterol

Appendix 5: Co-morbidity

	ELEVANCE
	ardiomyopathy is seen with cirrhosis and alcohol. Several liver-lung syndromes (HPS/PPHT/HHT) are described. A
	nical history, contrast echocardiography and other invasive tests may be necessary.
	ny history of cerebrovascular disease should be sought and described i.e. haemorrhagic/embolic/infarct and
	levant imaging and investigations supplied.
	I respiratory disease and investigations should be described, including any ITU admissions and performance
	atus. For hypoxic patient, include cross-sectional imaging, lung function, transfer studies and tests looking for
PE	Es/Shunts & AVMs
	or patients with chronic kidney disease (CKD) include investigations and treatments. If the eGFR
	30mls/min/1.73m ² for more than 3 months, a combined liver kidney graft may be necessary. Urinalysis, Albumin to
	reatinine ration (ACR), renal ultrasound and cultures should be completed in all patients with an eGFR
	60mls/min/1.73m ² .
	ssess the FRAX score in all patients (https://www.shef.ac.uk/FRAX/tool.jsp)
	V on treatment, is not a contraindication to liver transplantation. Enclose details from specialist teams of treatment
•	story, adherence and drug prescription.
-	ecord BMI and estimate dry BMI if ascites present. BMI>40kg/m ² is a relative contraindication
9	I surgical history should be detailed in the referral letter
	dietetic assessment prior to referral is mandatory. Offer appropriate nutritional support (e.g. protein supplements
	nd/or NGT feeding)
	equest an HbA1c (note anHbA1c may be artificially low in patients with chronic anaemia)
	or diabetes, document the duration, time on treatment (including years on insulin therapy), urinalysis, eye
	ramination & fundoscopy, vascular complications.
	ecord other components of metabolic syndrome such as hypertension and dyslipidaemia. history of cancer may not be a contraindication
	etails of dates of diagnosis, staging, treatment, prognosis, 5 year survival and correspondence from treating
	icologist
	ny communicable disease (including TB) in the patient and household contacts should be reported
	busing, next of kin, adherence with appointments, tests and therapies should be included in the referral letter
	earning disabilities or deafness/ visual impairment does not prevent assessment for transplantation
	cohol, illicit and prescribed drug usage including analgesics and sedatives, must be reported
abuse	series, musicant processed and dought mondaring analysis and socialities, musicas reported
Mental health Pri	ior history and treatment for mental illness, including self-harm should be detailed in the referral letter
Smoking Ac	ctive smokers should be advised to stop and be referred to smoking cessation service