

<b>Univariate regression of association with CRC</b>			
<b>Factor</b>	<b>Odds ratio</b>	<b>95% CI</b>	<b>P value</b>
FIT $\geq$ 10 $\mu$ gHb/g	2.588	0.9860 to 8.118	0.0713
Fit > 60 $\mu$ gHb/g	3.454	0.9905 to 12.56	0.0499
FIT > 100 $\mu$ gHb/g	4.412	1.257 to 16.16	0.0195
Age	1.044	1.005 to 1.090	0.0383
Male sex	3.947	1.542 to 12.13	0.0077
Change in bowel habit	0.8360	0.3552 to 1.949	0.6763
IDA	1.551	0.6310 to 3.634	0.3191
Rectal bleeding	2.249	0.9515 to 5.270	0.0602
Abdominal pain	1.320	0.5193 to 3.123	0.5386
Weight loss	2.200	0.8598 to 5.250	0.0832
Number of symptoms	1.663	1.080 to 2.538	0.0186
Rectal bleeding + change in bowel habit	2.204	0.7690 to 5.556	0.1109
Rectal bleeding + IDA	2.714	0.4095 to 10.59	0.2053
Rectal bleeding + abdominal pain	3.812	1.307 to 9.849	0.0083
Rectal bleeding + weight loss	4.904	1.317 to 14.86	0.0084
IDA + change in bowel habit	1.910	0.2923 to 7.198	0.4047
IDA + abdominal pain	6.254	0.8809 to 29.10	0.0304
IDA + weight loss	4.138	0.6061 to 17.37	0.0808
<b>Multivariate regression of association with CRC</b>			
<b>Factors</b>	<b>Odds ratio</b>	<b>95% confidence interval</b>	<b>P value</b>
FIT >100 $\mu$ gHb/g	4.495	1.224 to 17.45	0.0231
Rectal bleeding + weight loss	8.366	1.484 to 41.39	0.0098

**Supplemental table 3** - Univariate and multivariate regression for association with CRC